

Stroke

American Stroke
AssociationSM

JOURNAL OF THE AMERICAN HEART ASSOCIATION

A Division of American
Heart Association



Noninvasive Brain Stimulation May Improve Stroke-Related Dysphagia: A Pilot Study

Sandeep Kumar, Cynthia W. Wagner, Colleen Frayne, Lin Zhu, Magdy Selim, Wuwei Feng and Gottfried Schlaug

Stroke 2011;42;1035-1040; originally published online Mar 24, 2011;

DOI: 10.1161/STROKEAHA.110.602128

Stroke is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214
Copyright © 2011 American Heart Association. All rights reserved. Print ISSN: 0039-2499. Online
ISSN: 1524-4628

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://stroke.ahajournals.org/cgi/content/full/42/4/1035>

Subscriptions: Information about subscribing to Stroke is online at
<http://stroke.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters
Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax:
410-528-8550. E-mail:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/reprints>

Noninvasive Brain Stimulation May Improve Stroke-Related Dysphagia

A Pilot Study

Sandeep Kumar, MD; Cynthia W. Wagner, MS, CCC-SLP; Colleen Frayne, MS, CCC-SLP; Lin Zhu, BS; Magdy Selim, MD, PhD; Wuwei Feng, MD, MS; Gottfried Schlaug, MD, PhD

Background and Purpose—Treatment options for stroke-related dysphagia are currently limited. In this study, we investigated whether noninvasive brain stimulation in combination with swallowing maneuvers facilitates swallowing recovery in dysphagic stroke patients during early stroke convalescence.

Methods—Fourteen patients with subacute unilateral hemispheric infarction were randomized to anodal transcranial direct current stimulation (tDCS) versus sham stimulation to the sensorimotor cortical representation of swallowing in the unaffected hemisphere over the course of 5 consecutive days with concurrent standardized swallowing maneuvers. Severity of dysphagia was measured using a validated swallowing scale, Dysphagia Outcome and Severity scale, before the first and after the last session of tDCS or sham. The effect of tDCS was analyzed in a multivariate linear regression model using changes in Dysphagia Outcome and Severity Scale as the outcome variable after adjusting for the effects of other potential confounding variables such as the National Institutes of Health Stroke Scale and Dysphagia Outcome and Severity scale scores at baseline, acute ischemic lesion volumes, patient age, and time from stroke onset to stimulation.

Results—Patients who received anodal tDCS gained 2.60 points of improvement in Dysphagia Outcome and Severity scale scores compared to patients in the sham stimulation group who showed an improvement of 1.25 points ($P=0.019$) after controlling for the effects of other aforementioned variables. Six out of 7 (86%) patients in tDCS stimulation group gained at least 2 points of improvement compared with 3 out of 7 (43%) patients in the sham group ($P=0.107$).

Conclusions—Because brain stem swallowing centers have bilateral cortical innervations, measures that enhance cortical input and sensorimotor control of brain stem swallowing may be beneficial for dysphagia recovery. (*Stroke*. 2011;42:1035-1040.)

Key Words: dysphagia ■ noninvasive brain stimulation ■ stroke recovery ■ swallowing recovery
■ transcranial direct current stimulation

Dysphagia is a potentially fatal complication of stroke.¹ It afflicts numerous patients with hemispheric strokes¹ and has high rates of complications, even after adjusting for stroke severity.² Because hemispheric infarcts are the major subtype of ischemic stroke in the population,³ it can be assumed that the magnitude of dysphagia burden attributable to such strokes is large. Despite its frequent occurrence, treatment of stroke-related dysphagia remains limited. The usual practice is to provide nutritional support via alternative feeding methods, until swallowing functions recover; however, such methods fail to protect against complications of dysphagia such as aspiration pneumonia.^{4,5} Development of an effective intervention that improves swallowing in the early course of stroke recovery will be helpful in curtailing dysphagia-related complications and improving swallowing functions.

Swallowing functions are subserved by a distributed brain network, although involvement of the inferior peri-rolandic sensorimotor cortex appears consistent across studies.^{6–8} Disruption of projections from these cortical regions to the brain stem “swallowing centers” produces dysphagia with hemispheric strokes.⁹ Different lines of evidence suggest that recovery of swallowing functions occurs via expansion of the pharyngeal representation in the uninvolved hemisphere, possibly ensuring greater input to the brain stem swallowing centers.^{10,11} Cortical stimulation techniques may facilitate this process in patients with hemispheric lesions, in whom the brain stem and peripheral structures are intact but the upper echelons of the swallowing apparatus are dysfunctional. Repetitive transcranial magnetic stimulation over the swallowing motor cortex in healthy volunteers induces a long-term effect on the excitability of corticobulbar projections to

Received September 9, 2010; accepted November 19, 2010.

From the Department of Neurology (S.K., L.Z., M.S., W.F., G.W.), Stroke Division, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA; Voice, Speech, and Swallowing Service (C.W.W., C.F.), Beth Israel Deaconess Medical Center, Boston, MA.

Correspondence to Sandeep Kumar, MD, Beth Israel Deaconess Medical Center and Harvard Medical School, 330 Brookline Avenue, Palmer 127 Boston, MA 02215. E-mail skumar@bidmc.harvard.edu

© 2011 American Heart Association, Inc.

Stroke is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.110.602128

the pharynx¹² and may improve swallowing functions in dysphagic stroke patients.¹³

Transcranial direct current stimulation (tDCS) is another noninvasive brain stimulation technique that utilizes weak direct current to produce shifts in neuronal excitability^{14,15} and can be combined with swallowing maneuvers or exercises. It has generated great interest recently for its ease of use, patient tolerability, and safety profile, which is of particular importance during the acute/subacute phases of a stroke. It has been shown to improve motor functions in chronic stroke patients.^{16,17} Moreover, presence of a sham mode makes it possible to examine its effects in a blinded trial paradigm.¹⁸ More recently, investigators have shown that application of anodal tDCS to the pharyngeal motor cortex in healthy human subjects increases pharyngeal excitability in an intensity-dependent manner.¹⁹ In this pilot study, we investigated the effects of anodal tDCS versus sham stimulation of the unaffected hemisphere for improving dysphagia in the acute–subacute stroke phase.

Materials and Methods

This was an investigator-initiated, prospective, single-center, blinded pilot trial. All participants were recruited from our inpatient stroke service, were between 24 to 168 hours after their first ischemic stroke at time of enrollment, and had dysphagia secondary to a new unilateral hemispheric infarction. They were all evaluated by speech and language pathologists specializing in dysphagia (C.W. and C.F.) who were blinded to study allocation and rated swallowing impairments using a validated dysphagia scale, Dysphagia Outcome and Severity scale (DOSS).²⁰ DOSS scores range from 1 to 7, with 7 representing normal swallowing and 1 representing severe dysphagia. DOSS rates the functional severity of dysphagia and recommends a dietary level, independence level, and type of nutrition based on the level of impairment, thus conveying information about dysphagia severity and related disability. To qualify, a DOSS score of ≤ 5 (mild–severe dysphagia) was required. Patients with difficulty following instructions because of obtundation or cognitive impairment, preexisting swallowing problems, or other contraindications to tDCS were excluded.

All swallowing evaluations were conducted using hospital-based protocols that used different food consistencies representing the range of food consistencies consumed in real life (teaspoon, cup sip and straw sip of thin liquids, nectar, and thick liquids; honey; pureed solids; and a cookie). Patients were monitored for bolus control, oropharyngeal delays and retention, overt signs of aspiration including coughing, change in voice quality, or oxygen desaturation, with each consistency. In cases of ambiguity about assigning an appropriate DOSS score, a video swallow evaluation using the following boluses was performed the same day: teaspoon (3 mL) of nectar/thick liquid once, cup sip of nectar/thick liquid once (10 mL), straw sip of nectar/thick liquid once, followed by a teaspoon (3 mL) of thin liquid twice, cup sip (10 mL) followed by straw sips of thin liquid twice, followed by 5 mL Varibar pudding twice and half a vanilla wafer cookie twice. Overall, 7 patients required a video-fluoroscopic swallowing evaluation to record DOSS scores.

We recorded patient age, gender, lesion site, time in hours from stroke onset (time when patient was last seen normal if precise time of onset was unknown) to stimulation, lesion volume, and NIHSS scores as measures of stroke severity before stimulation. Acute ischemic lesion volumes were computed on diffusion-weighted imaging sequences on patient's brain MRI using customized software routines. The details of specific MRI sequence parameters, imaging processing, and volumetric analysis are described elsewhere.²¹ Two patients, unable to undergo an MRI, had their ischemic lesion volumes computed on a subacute head CT (obtained within 48–96 hours after symptoms onset). Patients were randomized to




Figure. Coregistration of anode mid-distance between C3/T3 using 10- to 20-EEG systems with a T1-weighted brain MRI demonstrates it to be centered over the caudal end of the primary motor cortex in a healthy volunteer.

receive either anodal tDCS or sham stimulation to the unaffected hemisphere using simple randomization and were blinded to their stimulation allocations. Using the international 10- to 20-EEG electrode system for guidance,²² a saline-soaked anodal electrode was placed over the undamaged hemisphere, mid-distance between C3 and T3 on the left or C4 and T4 on the right, with a reference electrode over the contralateral supraorbital region. This montage was expected to generate maximal current density over the inferior sensorimotor cortex and the neighboring premotor brain regions critical for reorganization of the swallowing motor cortex after a dysphagic stroke.^{10,11,23} We confirmed the location of the stimulating electrode and its proximity to the targeted regions by coregistering it with high-resolution T1-weighted MRI scans (Figure). A DOSS score was obtained immediately before stimulation sessions (DOSS-pre) and after the fifth session (DOSS-post).

The tDCS/sham was applied in conjunction with standardized swallowing maneuvers to provide adequate sensory and motor activation of the swallowing cortex.²⁴ All participants sucked on a lemon-flavored lollipop during these sessions. Patients reporting dryness of mouth were provided with 1 to 2 small ice chips intermittently. Patients were instructed to “swallow hard” every 30 seconds, thereby generating approximately 60 effortful swallows during each session. We used gesticulations to encourage aphasic patients to swallow at regular intervals. Occurrence of a swallow response was assessed by observing the movement of the thyroid cartilage or by palpating its excursion in patients with thicker necks. All subjects were able to follow study swallowing instructions appropriately. Anodal tDCS (2 mA for 30 minutes) or sham was

Table 1. Clinical Radiological Characteristics and Swallowing Measures of Enrolled Patients

Patient	Age	Sex	NIHSS Score	NIHSS Subscale Score (1a; 1)	Infarct Location	Lesion Volume (mL)	Time to Stimulation (Hour)	Dietary Status (Baseline)	DOSS-Pre	DOSS-Post		
						Anodal	tDCS	Group				
1	92	M	6	0; 1	Frontal, parietal, temporal lobes	20.6	40	NPO	1	5		
2	81	F	21	1; 2	Basal ganglia, internal capsule, parietal lobe	122.2	82	NPO	1	3		
3	90	F	10	0; 1	Insula, frontal lobe	43.9	50	Nectar-thick and pureed solids only	3	5		
4	70	F	9	0; 2	Insula, frontal lobe	36.48	30	NPO	1	6		
5	77	F	17	1; 2	Insula, frontal lobe, basal ganglia, internal capsule	58.06	97	Thin liquids, pureed solids	5	7		
6	84	M	20	1; 4	Internal capsule, frontal, temporal, parietal lobes	120	140	NPO	1	2		
7	64	M	12	1; 0	Basal ganglia, internal capsule	20.1	123	Nectar thick liquids, pureed solids	3	5		
						79.7*	13.6*	0.6*; 1.7*	60.2*	80.3*	2.1*	4.7*
						Sham	Group					
1	57	F	16	0; 2	Insula, frontal lobe, basal ganglia	84.65	42	Nectar thick and pureed solids	4	5		
2	83	M	12	0; 1	Insula, frontal lobe, basal ganglia	63.06	52	Ground solids and nectar thick	3	4		
3	50	M	16	1; 5	Insula, frontal, temporal, parietal lobe, basal ganglia	135.24	75	NPO	1	3		
4	74	M	6	0; 0	Insula, frontal lobe	22.5	76	Nectar thick and pureed solids	3	5		
5	72	M	11	1; 1	Insula, basal ganglia, internal capsule	40.32	146	NPO	1	2		
6	78	F	15	1; 2	Insula, basal ganglia, internal capsule	54.9	148	NPO	1	3		
7	76	F	16	0; 2	Insula, basal ganglia, internal capsule	84.56	138	Nectar thick and ground solids	4	4		
						70*	13.14*	0.4*; 2*	69.46*	96.71*	2.4*	3.7*

DOSS-post indicates Dysphagia Outcome and Severity scale after fifth stimulation session; DOSS-pre, Dysphagia Outcome and Severity scale before stimulation sessions; NIHSS, National Institutes of Health Stroke scale; NPO, nothing by mouth; tDCS, transcranial direct current stimulation.

*Average values for each column.

applied daily to the nonlesional hemisphere for 5 consecutive days. The tDCS was delivered through a battery-driven constant current stimulator (Phoresor; Iomed, Salt Lake City, UT), with the following electrode dimensions: 3×5 cm for the anode and 5×6 cm for the reference electrode.

The study was approved by our Institutional Review Board. A written informed consent was obtained from the patients or their legal representative before enrollment.

Statistical Analysis

We analyzed the effect of stimulation (tDCS or sham, entered as a binary variable) on improvement in dysphagia scores after adjusting for the potential confounding effects of other important variables, ie, stroke severity as assessed by baseline NIHSS score, ischemic lesion volume, baseline DOSS score, patient age, and time from stroke onset to stimulation. A correlation analysis and collinearity assessment among all independent variables were checked before the final model. A responder variable (yes or no) was defined as at least ≥2 points of improvement on DOSS. A logistic regression was applied with the same covariates from the general linear regression model as

a secondary outcome analysis. All statistical analyses were performed using SAS version 9.1 (SAS Institute, Cary, NC).

Results

Fourteen patients were recruited and randomized to anodal tDCS or sham stimulation group in a 1:1 fashion. The important characteristic of our patient sample is tabulated in Table 1. All patients who consented to participate in this study tolerated the sessions well; stimulation was not curtailed in anyone because of discomfort or fatigue. No adverse events, such as seizures, headaches, visual disturbances, or significant skin irritation, were observed. Two patients in the sham group but none in the tDCS group underwent percutaneous endoscopic gastrostomy placement after their trial participation.

Multivariate Analysis

NIHSS scores and DOSS scores at baseline, acute ischemic lesion volume, time to stimulation, and age were initially

Table 2. Results of Anodal tDCS Versus Sham Stimulation in Multivariate Analysis

	Anodal tDCS	Sham	<i>P</i>
Change in DOSS scores	2.6‡ (1.91–3.29)	1.26‡ (0.57, 1.95)	0.019†
≥2-point improvement in DOSS score	6/7 (86%)	3/7 (43%)	0.107*

**P* is based on a logistic regression model with baseline DOSS, NIHSS, age, time to treatment, and stimulation group as covariates.

†*P* based on a general linear regression model with baseline DOSS, NIHSS, age, time to treatment, and stimulation group as covariates.

‡Least square mean and 95% confidence interval estimated from the general linear model.

included for a generalized linear model; however, NIHSS and lesion volume were highly correlated ($r=0.84$ and $P=0.0002$), and further collinearity diagnostics revealed a significant collinearity (tolerance >0.1 ; variance inflation factor <10) between the 2 variables. Thus, the latter was eliminated from the model. In summary (Table 2), our results show that patients who received anodal tDCS gained 2.60 (95% CI, 1.91–3.29) points on DOSS, whereas patients in the sham stimulation group improved by 1.25 (95% CI, 0.57–1.95); the difference between 2 groups reached a statistical significance with $P=0.019$. DOSS at baseline ($P=0.045$) and NIHSS at baseline ($P=0.049$) were significantly associated with improvement on DOSS scores. Age ($P=0.228$) was not a good predictor for improvement based on our model analysis. Our secondary outcome included a logistic regression analysis that was based on at least 2 points of improvement with DOSS. Six out of 7 patients (86%) in the tDCS group had ≥ 2 points of improvement on their DOSS scores versus 3 out of 7 patients (43%) in the sham stimulation group ($P=0.107$).

Discussion

The findings of this pilot study show that repeated application of anodal tDCS to the unaffected swallowing cortex in combination with timed effortful swallowing is associated with significant swallowing improvement over sham after adjusting for the effects of baseline stroke and dysphagia severity, age, and time to stimulation in patients with acute–subacute unilateral hemispheric infarction. Our results also attest to the feasibility and tolerability of tDCS in this stroke subpopulation during early phases of stroke recovery.

The brain stimulation effect might be explained by an augmentation effect of the naturally occurring changes in the unaffected swallowing cortex.^{10,11} Combining the sensorimotor effects of swallowing maneuvers with simultaneous brain stimulation of the unaffected hemisphere may have been an important component of the effect. Sensory input from the pharynx is known to increase excitability of the swallowing sensorimotor cortex through convergent afferent activity,¹² and pharyngeal sensory stimulation in dysphagic stroke patients produces an increase in the excitability of the swallowing motor cortex of the unaffected hemisphere.²⁴ However, studies investigating induction of plasticity in the human motor cortex using paired associative paradigm have

shown that cortical stimulation, if paired with peripheral stimulation of the somatosensory afferents, leads to greater increases in cortical excitability than produced by stimulation alone and induces topographically specific plastic changes.²⁵ This increase in excitability was prevented by using dextromethorphan, which is known to block development of long-term-potential.²⁶ In animal studies, motor skill learning has been shown to produce long-term potentiation and long-term depression, leading to changes in synaptic strength in the primary motor cortex.²⁷ Cortical stimulation studies in experimental stroke models have shown stronger effects when peripheral sensorimotor activities were combined with central stimulation.²⁸ More recently, investigators²⁹ have shown that training in humans or low-frequency stimulation in mouse M1 slices produces release of brain-derived neurotrophic factor, which is necessary to induce long-term synaptic plasticity from direct current stimulation. In chronic stroke patients, combining peripheral nerve stimulation or peripheral sensorimotor activities with tDCS facilitates the beneficial effects of training on motor performance beyond levels reached by each intervention alone.^{30,31} Thus, data from diverse sources indicate that combining repetitive peripheral sensorimotor stimulation with noninvasive brain stimulation can potentiate relearning and consolidation of motor skills to a level unattainable by any of these interventions alone in subacute or chronic stroke patients and appears to have benefitted our subjects.

Our statistical methods were designed to control for discrepancies of important predictors of dysphagia recovery between groups that the randomization procedures may have failed to correct in our small sample. Although there is little data published on predictors of dysphagia recovery in stroke patients, baseline NIHSS score,³² stroke lesion volume,³³ and age³⁴ have been found to be important factors influencing functional recovery in stroke patients and were included in the analysis. Because swallowing functions in our patients were expected to recover over time, time to stimulation was also included in our analysis. Our model shows that baseline NIHSS, DOSS scores, and anodal tDCS were associated with improvement. Introduction of all these variables could have overfitted our model and exhausted degrees of freedom for estimation with a small sample size. However, the intent of this analysis was to gain an understanding about the important covariates influencing swallowing recovery and adjust for their effects on experimental treatment and not to try to build a predictive model.

It is possible that in a minority of patients, especially in those with more circumscribed lesions, the ipsilesional hemisphere may have played a role in swallowing recovery and accounted for some variability in responses to stimulation. This poses an important question whether uniform application of anodal tDCS to the uninvolved hemisphere will benefit all such patients. However, because brain stem swallowing centers have bilateral innervations with little evidence for transcallosal inhibition,³⁵ we hypothesized that stimulation of either hemisphere would produce an increase in pharyngeal excitability. Furthermore, stimulation of the uninvolved hemisphere was less likely to be affected by neuronal loss or tissue damage and responses would be more

uniform; stimulating the nonlesioned hemisphere was also expected to be safer with respect to any potential seizures risk or tissue damage in the acute stroke phase. The optimal dose for stimulating the pharyngeal motor cortex has not been established. A recent report suggests that doses higher than that used for stimulating the primary motor cortex are necessary to produce comparable responses from the swallowing cortex.¹⁹ Our protocol predates the publication of this report and alternative doses can be tried in future studies to assess their superiority. We chose our dose based on previous study protocols that have shown that application of 2 mA to the dorsolateral frontal lobes is effective and well-tolerated.³⁶ Our decision to perform 5 sessions of stimulation was based on recent reports showing an additive effect of repeated session of tDCS³⁷ and taking logistical considerations such as duration of hospitalization in mind. It is possible that more sessions may have produced a stronger effect. Other study limitations include nonroutine use of video-fluoroscopic swallowing evaluations in all subjects, which were performed based on clinical judgment of evaluating speech and language pathologists. Although DOSS has excellent inter-rater reliability,²⁰ our methods of evaluation may have failed to account for some random variability in assigning DOSS scores in this study. In addition, use of a single evaluation scheme for determining swallowing functions may have been unable to capture pertinent details about changes in swallowing physiology in these subjects. In future studies, additional dysphagia assessment scoring tools should be obtained to tests the robustness of any treatment effect.

Conclusions

In conclusion, the results from this pilot study show a promising efficacy of anodal tDCS application with the swallowing cortex of the unaffected hemisphere combined with effortful swallowing maneuvers for improving dysphagia in stroke patients. Further studies are warranted to refine this promising intervention by exploring effects of stimulation parameters, frequency of stimulation, and timing of the intervention in improving swallowing functions in dysphagic stroke patients.

Acknowledgments

The authors thank Drs Vijay Renga and Dinesh Nair for helping with the noninvasive brain stimulation of our initial patients.

Sources of Funding

Dr Kumar receives partial salary support from the NIH (NINDS 5U01-NS044876-03 [Site PI]) and receives research support from the Charles and Irene Goldman Neurology Research Fund (PI). Dr Selim receives research support from the NIH (NINDS 1R01-NS 057127-01A1 [PI], NINDS 1R01-NS 045754-01A2 [Co-I], and 5R01-HL46690-14 [Co-I]). Dr Schlaug receives research support from the NIH (NIDCD 1R01 DC008796 [PI], NIDCD 3R01DC008796-02S1 [PI], R01 DC009823-01 [PI], and 1R01-NS 057127 [Co-I]). This study received funding support from NIH (R01 DC008796 and R01 DC009823 to Dr Schlaug [PI] and CIMIT to Dr Schlaug [PI]).

Disclosures

None.

References

- Martino R, Foley N, Bhogal S, Diamant N, Speechley M, Teasel R. Dysphagia after stroke. Incidence, diagnosis, and pulmonary complications. *Stroke*. 2005;36:2756–2763.
- Barer DH. The natural history and functional consequences of dysphagia after hemispheric stroke. *J Neurol Neurosurg Psychiatry*. 1989;52:236–241.
- Turney TM, Garraway WM, Whisnant JP. The natural history of hemispheric and brainstem infarction in Rochester, Minnesota. *Stroke* 1984; 15:790–794.
- Langmore SE, Terpenning MS, Schork A, Chen Y, Murray JT, Lopatin D, Loesche WJ. Predictors of aspiration pneumonia: how important is dysphagia? *Dysphagia*. 1998;13:69–81.
- Dziewas R, Ritter M, Schilling M, Konrad C, Oelenberg S, Nabavi DG, Stögbauer F, Ringelstein EB, Lüdemann P. Pneumonia in acute stroke patients fed by nasogastric tube. *J Neurol Neurosurg Psychiatry*. 2004; 75:852–856.
- Penfield W, Boldery E. Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain*. 1937; 60:389–443.
- Martin RE, Sessle BJ. The role of the cerebral cortex in swallowing. *Dysphagia*. 1993;8:195–202.
- Martin RE, Goodyear BG, Gati JS, Menon RS. Cerebral cortical representation of automatic and volitional swallowing in humans. *J Neurophysiol*. 2001;85:938–950.
- Hamdy S, Aziz Q, Rothwell JC, Crone R, Hughes D, Tallis RC, Thompson DG. Explaining oropharyngeal dysphagia after unilateral hemispheric stroke. *Lancet*. 1997;350:686–692.
- Hamdy S, Aziz Q, Rothwell JC, Power M, Singh KD, Nicholson DA, Tallis RC, Thompson DG. Recovery of swallowing after dysphagic stroke relates to functional reorganization in the intact motor cortex. *Gastroenterology*. 1998; 115:1104–1112.
- Li S, Luo C, Yu B, Yan B, Gong Q, He C, He L, Huang X, Yao D, Lui S, Tang H, Chen Q, Zeng Y, Zhou D. Functional magnetic resonance imaging study on dysphagia after unilateral hemispheric stroke: a preliminary study. *J Neurol Neurosurg Psychiatry*. 2009;80:1320–1329.
- Gow D, Rothwell J, Hobson A, Thompson D, Hamdy S. Induction of long-term plasticity in human swallowing cortex following repetitive cortical stimulation. *Clin Neurophysiology*. 2004;115:1044–1051.
- Khedr EM, Abo-Elfetoh N, Rothwell JC. Treatment of post-stroke dysphagia with repetitive transcranial magnetic stimulation. *Acta Neurol Scand*. 2009;119:155–161.
- Nitsche MA, Paulus W. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *J Physiol*. 2000; 527:633–639.
- Nitsche MA, Paulus W. Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans. *Neurology*. 2001; 57:1889–1901.
- Hummel F, Celnik P, Giraux P, Floel A, Wu WH, Gerloff C, Cohen LG. Effects of non-invasive cortical stimulation on skilled motor function in chronic stroke. *Brain*. 2005;128:490–499.
- Schlaug G, Renga V, Nair D. Transcranial Direct Current Stimulation in Stroke Recovery. *Arch Neurol*. 2008;65:1–7.
- Gandiga PC, Hummel FC, Cohen LG. Transcranial DC stimulation (tDCS): a tool for double-blind sham-controlled clinical studies in brain stimulation. *Clin Neurophysiol*. 2006;117:845–850.
- Jefferson S, Mistry S, Singh S, Rothwell J, Hamdy S. Characterizing the application of transcranial direct current stimulation in human pharyngeal motor cortex. *Am J Physiol Gastrointest Liver Physiol*. 2009;297: G1035–G1040.
- O'Neil KH, Purdy M, Falk J, Gallo L. The Dysphagia Outcome and Severity Scale. *Dysphagia*. 1999;14:139–145.
- Schlaug G, Benfield A, Baird AE, Siewert B, Lövblad KO, Parker RA, Edelman RR, Warach S. The ischemic penumbra: operationally defined by diffusion and perfusion MRI. *Neurology*. 1999;53:1528–1537.
- Steinmetz H, Gunter F, Bernd-Ulrich M. Craniocerebral topography within the international 10–20 system. *Electroencephalogr Clin Neurophysiol*. 1989;72:499–506.

23. Miranda PC, Lomarev M, Hallett M. Modeling the current distribution during transcranial direct current stimulation. *Clin Neurophysiol.* 2006;117:1623–1629.
24. Fraser C, Power M, Hamdy S, Rothwell J, Hobday D, Hollander I, Tyrell P, Hobson A, Williams S, Thompson D. Driving plasticity in human adult motor cortex is associated with improved motor function after brain injury. *Neuron.* 2002;34:831–840.
25. Stefan K, Kunesch E, Cohen LG, Benecke R, Classen J. Induction of plasticity in the human motor cortex by paired associative stimulation. *Brain.* 2000;123:572–584.
26. Stefan K, Kunesch E, Benecke R, Cohen LG, Classen J. Mechanisms of enhancement of human motor cortex excitability induced by interventional paired associative stimulation. *J Physiol.* 2002;543:699–708.
27. Rioult-Pedotti MS, Friedman D, Donoghue JP. Learning-induced LTP in neocortex. *Science.* 2000;290:533–536.
28. Adkins-Muir DL, Jones TA. Cortical electrical stimulation combined with rehabilitative training: enhanced functional recovery and dendritic plasticity following focal cortical ischemia in rats. *Neurol Res.* 2003;25:780–788.
29. Fritsch B, Reis J, Martinowich K, Schambra HM, Ji Y, Cohen LG, Lu B. Direct current stimulation promotes BDNF-dependent synaptic plasticity: potential implications for motor learning. *Neuron.* 2010;66:198–204.
30. Celnik P, Paik NJ, Vandermeeren Y, Dimyan M, Cohen LG. Effects of combined peripheral nerve stimulation and brain polarization on performance of a motor sequence task after chronic stroke. *Stroke.* 2009;40:1764–1771.
31. Lindenberg R, Renga V, Zhu LL, Nair D, Schlaug G. Bihemispheric brain stimulation facilitates motor recovery in chronic stroke patients. *Neurology.* 2010;75:2176–2184.
32. Adams HP Jr, Davis PH, Leira EC, Chang KC, Bendixen BH, Clarke WR, Woolson RF, Hansen MD. Baseline NIH Stroke Scale score strongly predicts outcome after stroke: A report of the Trial of Org 10172 in Acute Stroke Treatment (TOAST). *Neurology.* 1999;53:126–131.
33. Thijs VN, Lansberg MG, Beaulieu C, Marks MP, Moseley ME, Albers GW. Is early ischemic lesion volume on diffusion-weighted imaging an independent predictor of stroke outcome? A multivariable analysis. *Stroke.* 2000;31:2597–2602.
34. Turhan N, Atalay A, Muderrisoglu H. Predictors of functional outcome in first-ever ischemic stroke: A special interest to ischemic subtypes, comorbidity and age. *NeuroRehabilitation.* 2009;24:321–326.
35. Hamdy S, Aziz Q, Rothwell JC, Hobson A, Thompson DG. Sensorimotor modulation of human cortical swallowing pathways. *J Physiol.* 1998;506:857–866.
36. Iyer MB, Mattu U, Grafman J, Lomarev M, Sato S, Wassermann EM. Safety and cognitive effect of frontal DC polarization in healthy individuals. *Neurology.* 2005;64:872–875.
37. Boggio PS, Nunes A, Rigonatti SP, Nitsche MA, Pascual-Leone A, Fregni F. Repeated sessions of noninvasive brain DC stimulation is associated with motor function improvement in stroke patients. *Rest Neurol Neurosci.* 2007;25:123–129.